



HIGHLIGHTS OF PRESCRIBING INFORMATION	
These highlights do not include all the information needed to use ARGATROBAN INJECTION safely and effectively. See full prescribing information for ARGATROBAN INJECTION.	
ARGATROBAN INJECTION, for intravenous infusion only	
Initial U.S. Approval: 2000	
—INDICATIONS AND USAGE—	
Argatroban is a direct thrombin inhibitor indicated:	
• For prophylaxis or treatment of thrombosis in adult patients with heparin-induced thrombocytopenia (HIT) (1, 1)	
• As an anticoagulant in adults patients with or at risk for HIT undergoing percutaneous coronary intervention (PCI) (1, 2)	
—DOSAGE AND ADMINISTRATION—	
• Argatroban Injection 50 mg/50 mL (1 mg/mL) is ready for intravenous infusion. Dilution is not needed. (2, 2)	
Heparin-Induced Thrombocytopenia	
The dose for heparin-induced thrombocytopenia without hepatic impairment is 2 mg/kg/min administered as a continuous infusion. (2, 2)	
Percutaneous Coronary Intervention	
The dose for patients with or at risk for heparin-induced thrombocytopenia undergoing percutaneous coronary intervention is started at 0.2 mg/kg/min and a bolus of 350 mg/kg administered via a large bore intravenous line over 3 to 5 minutes. (2, 3)	
DOSAGE FORMS AND STRENGTHS	
• Injection: 50 mg/50 mL (1 mg/mL) ready for intravenous infusion single-dose vial. (3)	
—CONTRAINDICATIONS—	
• Major bleeding (4)	
• History of hypersensitivity to this product (4)	

WARNINGS AND PRECAUTIONS	
• Hemorrhage can occur. Unexplained fall in hematocrit or blood pressure may indicate hemorrhage (5, 1)	
• Hepatic impairment: Adjust starting dose and titrate carefully in patients with HT who have moderate or severe hepatic impairment. Avoid use in PCI in patients with clinically significant hepatic impairment. (5, 2)	
ADVERSE REACTIONS	
• HIT patients: The most common (>5%) adverse reactions were dyspnea, hypotension, fever, diarrhea, hsp, and cardiac arrest. (6, 1)	
• PCI patients: The most common (>5%) adverse reactions were chest pain, hypotension, back pain, nausea, vomiting and headache. (6, 1)	
To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.	
DRUG INTERACTIONS	
• Heparin: Allow sufficient time for heparin's effect on activated partial thromboplastin time (aPTT) to decrease before initiating Argatroban Injection therapy. (7, 1)	
• Warfarin: Concomitant use results in increased prolongation of PT and INR. (7, 2)	
• Thrombolytic agents or glycoprotein IIb/IIIa antagonists: Safety and effectiveness of concomitant use with argatroban have not been established. (7, 3, 4)	
USE IN SPECIFIC POPULATIONS	
• Lactation: Discontinue nursing or drug taking into account the importance of the drug to the mother. (8, 2)	
• Pediatric Use: Safety and effectiveness have not been established. (8, 4)	
See 17 for PATIENT COUNSELING INFORMATION	

Revised: 03/2023

FULL PRESCRIBING INFORMATION: CONTENTS*	
1 INDICATIONS AND USAGE	
1.1 Heparin-Induced Thrombocytopenia	
1.2 Percutaneous Coronary Intervention	
2 DOSAGE AND ADMINISTRATION	
2.1 Preparation for Intravenous Administration	
2.2 Dosing in Patients with Heparin-Induced Thrombocytopenia	
2.3 Dosing in Patients Undergoing Percutaneous Coronary Intervention	
2.4 Conversions to Oral Anticoagulant Therapy	
3 DOSAGE FORMS AND STRENGTHS	
4 CONTRAINDICATIONS	
5 WARNINGS AND PRECAUTIONS	
5.1 Risk of Hemorrhage	
5.2 Hepatic Impairment	
5.3 Laboratory Tests	
6 ADVERSE REACTIONS	
6.1 Hemorrhage	
6.2 Hepatic Impairment	
7 DRUG INTERACTIONS	
7.1 Heparin	
7.2 Oral Anticoagulant Agents	
7.3 Aspirin/Acetaminophen	
7.4 Thrombolytic Agents	
7.5 Glycoprotein IIb/IIIa Antagonists	

FULL PRESCRIBING INFORMATION	
1 INDICATIONS AND USAGE	
1.1 Heparin-Induced Thrombocytopenia	
Argatroban Injection is indicated for prophylaxis or treatment of thrombosis in adult patients with heparin-induced thrombocytopenia (HIT).	
1.2 Percutaneous Coronary Intervention	
Argatroban Injection is indicated as an anticoagulant in adult patients with or at risk for HIT undergoing percutaneous coronary intervention (PCI).	
2 DOSAGE AND ADMINISTRATION	
2.1 Preparation for Intravenous Administration	
Dilution is not required for Argatroban Injection 50 mg/50 mL (1 mg/mL).	
Argatroban Injection 50 mg/50 mL (1 mg/mL)	
Each 50 mL glass vial contains 50 mg argatroban (1 mg/mL), and, as supplied, is ready for intravenous infusion. Dilution is not required. Argatroban Injection is a clear, colorless to pale yellow solution. Parenteral drug products should be inspected for any particulate matter and discoloration prior to administration whenever solution and container permit. Do not use if solution is cloudy, contains precipitates, or if the flip of seals is not intact.	
Val may be inverted for use with a medical infusion set.	
2.2 Dosing in Patients with Heparin-Induced Thrombocytopenia	
For adult patients with HIT, discontinue heparin therapy and obtain a baseline activated partial thromboplastin time (aPTT). The recommended initial dose of argatroban for adult patients without hepatic impairment is 2 mg/kg/min, administered as a continuous infusion (see Table 1).	

Recommended Doses and Infusion Rates for 2 mg/kg/min Dose of Argatroban for Patients With HIT* and Without Hepatic Impairment (1 mg/mL Final Concentration)		
Body Weight (kg)	Dose (mg/min)	Infusion Rate (mL/hr)
50	100	6
60	120	7
70	140	8
80	160	10
90	180	11
100	200	12
110	220	13
120	240	14
130	260	16
140	280	17

*With or without thrombosis	
Monitoring Strategy	
For use in HIT, therapy with Argatroban Injection is monitored using the aPTT with a target range of 1.5 to 3 times the initial baseline value (not to exceed 100 seconds). Tests of anticoagulant effects (including the aPTT) typically attain steady-state levels within 1 to 3 hours following initiation of Argatroban Injection. Check the aPTT 2 hours after initiation of therapy and after every dose change to confirm that the patient has attained the desired therapeutic range.	
Dose Adjustment	
After the initiation of Argatroban Injection, adjust the dose (not to exceed 10 mg/kg/min) as necessary to obtain a steady-state aPTT in the target range (see <i>Clinical Studies</i> (14, 1)).	
2.3 Dosing in Patients Undergoing Percutaneous Coronary Intervention	
Initial Dosing	
Initiate an infusion of Argatroban Injection at 25 mg/kg/min and administer a bolus of 350 mg/kg via a large bore intravenous line over 3 to 5 minutes (see Table 2). Check an activated clotting time (ACT) 5 to 10 minutes after the bolus dose is completed. The PO procedure may proceed if the ACT is greater than 300 seconds.	
Dose Adjustment	
If the ACT is less than 300 seconds, an additional intravenous bolus dose of 150 mg/kg should be administered, the infusion rate decreased to 30 mg/kg/min, and the ACT checked 5 to 10 minutes later (see Table 2). If the ACT is greater than 450 seconds, decrease the infusion rate to 15 mg/kg/min, and check the ACT 5 to 10 minutes later (Table 3).	
Continue titrating the dose until a therapeutic ACT (between 300 and 450 seconds) has been achieved; continue the same infusion rate for the duration of the PCI procedure.	
In case of dissection, impending abrupt closure, thrombus formation during the procedure, or inability to open or maintain an ACT over 300 seconds, additional bolus doses of 150 mg/kg may be administered and the infusion dose increased to 40 mg/kg/min. Check the ACT after each additional bolus or change in the rate of infusion.	

Table 2 Recommended Starting and Maintenance Doses (Within the Target ACT Range) of Argatroban Injection in Patients Undergoing PCI Without Hepatic Impairment (1 mg/mL Final Concentration)				
Body Weight (kg)	Starting Bolus Dose (350 mcg/kg)		Starting and Maintenance Continuous Infusion Dosing For ACT 300-450 seconds 25 mcg/kg/min	
	Bolus Dose (mcg)	Bolus Volume (mL)	Continuous Infusion Dose (mcg/min)	Continuous Infusion Rate (mL/hr)
50	17500	18	1250	75
60	21000	21	1500	90
70	24500	25	1750	105
80	28000	28	2000	120
90	31500	32	2250	135
100	35000	35	2500	150
110	38500	39	2750	165
120	42000	42	3000	180
130	45500	46	3250	195
140	49000	49	3500	210

Table 3 Recommended Dose Adjustments of Argatroban Injection for Patients Outside of ACT Target Range Undergoing PCI Without Hepatic Impairment (1 mg/mL Final Concentration)						
Body Weight (kg)	If ACT Less than 300 seconds Dose Adjustment* 30 mcg/kg/min			If ACT Greater than 450 seconds Doseage Adjustment* 15 mcg/kg/min		
	Additional Bolus Dose (mg)	Bolus Volume (mL)	Continuous Infusion Dose (mg/min)	Continuous Infusion Rate (mL/hr)	Continuous Infusion Dose (mg/min)	Continuous Infusion Rate (mL/hr)
50	7500	8	1500	90	750	45
60	9000	9	1800	108	900	54
70	10500	11	2100	126	1050	63
80	12000	12	2400	144	1200	72
90	13500	14	2700	162	1350	81
100	15000	15	3000	180	1500	90
110	16500	17	3300	198	1650	99
120	18000	18	3600	216	1800	108
130	19500	20	3900	234	1950	117
140	21000	21	4200	252	2100	126

NOTE: 1 mg = 1000 mg; 1 kg = 2.2 lbs	
*No bolus dose is given if ACT greater than 450 seconds	
Monitoring Strategy	
For use in PCI, therapy with Argatroban Injection is monitored using the aPTT. After ACTs before dosing, 5 to 10 minutes after bolus dosing, following adjustments in the infusion rate, and at the end of the PCI procedure. Obtain additional ACTs every 20 to 30 minutes during prolonged procedure.	

Continued Anticoagulation after PCI	
If a patient requires anticoagulation after the procedure, Argatroban Injection may be continued, but at a rate of 1 mg/kg/min. If a patient requires anticoagulation after the procedure, Argatroban Injection may be continued, but at a rate of 1 mg/kg/min. If a patient requires anticoagulation after the procedure, Argatroban Injection may be continued, but at a rate of 1 mg/kg/min.	

Dosing in Patients with Hepatic Impairment

Initial Dosing

For adult patients with HT and moderate or severe hepatic impairment (based on Child-Pugh classification), an initial dose of 0.5 mg/kg/min is recommended, based on the approximately 4-fold decrease in argatroban clearance relative to those with normal hepatic function. Monitor the aPTT closely, and adjust the dosage as clinically indicated.

Monitoring Strategy

Achievement of steady-state aPTT levels may take longer and require more dose adjustments in patients with hepatic impairment compared to patients with normal hepatic function.

For patients with hepatic impairment undergoing PCI and who have HT or are at risk for HIT, carefully titrate argatroban until the desired level of anticoagulation is achieved. Achievement of steady-state aPTT levels may take longer and require more argatroban dose adjustments in patients with hepatic impairment compared to patients with normal hepatic function. In patients with clinically significant hepatic disease or AST/ALT levels ≥ 3 times the upper limit of normal should be avoided (see Warnings and Precautions (5.2)).

2.5 Conversion to Oral Anticoagulant Therapy

Initiating Oral Anticoagulant Therapy

When converting patients from argatroban to oral anticoagulant therapy, consider the potential for combined effects with argatroban with co-administration of argatroban and warfarin. A loading dose of warfarin should not be initiated. Initiation of oral anticoagulant therapy with warfarin should be initiated after argatroban is discontinued. In patients with hepatic impairment when initiating warfarin, it is suggested that Argatroban and warfarin therapy be overlapped. There are insufficient data available to recommend the duration of the overlap.

Co-administration of Warfarin and Argatroban Injection at Doses up to 2 mg/kg/min

Measure INR daily while argatroban and warfarin are co-administered. In general, with doses of argatroban injection up to 2 mg/kg/min, Argatroban Injection can be discontinued when the INR is > 4.0 combined with the desired therapeutic range. When the INR is > 4.0 and the desired therapeutic range is not achieved, resume the infusion of Argatroban Injection and repeat the procedure daily until the desired therapeutic range on warfarin alone is reached.

Co-administration of Warfarin and Argatroban Injection at Doses Greater than 2 mg/kg/min

For doses of argatroban injection greater than 2 mg/kg/min, the relationship of INR between warfarin alone to the INR on warfarin plus argatroban is less predictable. In this case, in order to predict the INR on warfarin alone, temporarily discontinue the dose of Argatroban Injection to a dose of 2 mg/kg/min. Repeat the INR on argatroban injection and warfarin. When the INR is > 4.0 and the desired therapeutic range is not achieved, resume the infusion of Argatroban Injection and repeat the procedure daily until the desired therapeutic range on warfarin alone is reached.

Discontinuation of Argatroban Injection at Doses up to 2 mg/kg/min

For doses of argatroban injection up to 2 mg/kg/min, the relationship of INR between warfarin alone to the INR on warfarin plus argatroban is less predictable. In this case, in order to predict the INR on warfarin alone, temporarily discontinue the dose of Argatroban Injection to a dose of 2 mg/kg/min. Repeat the INR on argatroban injection and warfarin. When the INR is > 4.0 and the desired therapeutic range is not achieved, resume the infusion of Argatroban Injection and repeat the procedure daily until the desired therapeutic range on warfarin alone is reached.

Discontinuation of Argatroban Injection at Doses Greater than 2 mg/kg/min

For doses of argatroban injection greater than 2 mg/kg/min, the relationship of INR between warfarin alone to the INR on warfarin plus argatroban is less predictable. In this case, in order to predict the INR on warfarin alone, temporarily discontinue the dose of Argatroban Injection to a dose of 2 mg/kg/min. Repeat the INR on argatroban injection and warfarin. When the INR is > 4.0 and the desired therapeutic range is not achieved, resume the infusion of Argatroban Injection and repeat the procedure daily until the desired therapeutic range on warfarin alone is reached.